

Variational Calculations for ^3He Impurities on ^4He Droplets

A. Belić^a, F. Dalfovo^b, S. Fantoni^c, and S. Stringari^b

^a *International Center for Theoretical Physics,*

P.O.Box 586, I-34100 Trieste, Italy

^b *Dipartimento di Fisica, Università di Trento,*

I-38050 Povo, Italy

^c *Interdisciplinary Laboratory, SISSA,*

Via Beirut 2, I-34014 Trieste, Italy

Abstract

Variational Monte Carlo method is used to calculate ground state properties of ^4He droplets, containing 70, 112, 168, 240, 330, and 728 particles. The resulting particle and kinetic energy densities are used as an input in the Feynman-Lekner theory for ^3He impurities. The kinetic energy density of ^4He atoms and the energy of the ^3He surface states are compared with the results of previous phenomenological calculations.

67.40

I. INTRODUCTION

Helium droplets have attracted some interest in recent years. A major motivation is the fact that they represent a prototype of finite size quantum systems, behaving as fluid of strongly interacting structureless particles. Several theoretical schemes have been developed for pure helium droplets [1–6]. The comparison between theory and experiments is still elusive, mainly because helium droplets are so weakly bound objects that their experimental characterization is very difficult. An interesting approach, from this point of view, is the use of atomic and molecular impurities as probes [7,8]. Theoretical calculations for droplets with impurities have been recently done with both Monte Carlo [9] and density functional methods [10,11]. Impurities heavier than helium atoms are expected to have small zero point motion, so that they can be treated as classical objects in a quantum fluid [11]. This is certainly not true for ^3He and hydrogen impurities, for which a full quantum mechanical treatment is needed. The case of ^3He impurities is particularly interesting from the theoretical viewpoint. Since ^3He and ^4He atoms interact through the same potential, the properties of their mixed systems are determined only by quantum effects, i.e., the different statistics and the different zero point motion. One ^3He atom, being lighter than ^4He , tends to move in regions of low ^4He density. This is the origin of the so called Andreev state of ^3He on a ^4He liquid-vapour interface at low temperature [12–17]. The same behavior is observed in ^4He films on solid substrates [18], where the layer structure of the ^4He density produces a rich variety of ^3He states [19–22]. Predictions on the Andreev states of ^3He atoms on ^4He droplets have been already given [10] using a phenomenological density functional, as well as the variational Feynman-Lekner theory. A key ingredient in the latter approach is the ^4He kinetic energy density, for which a simple approximate expression was proposed in Ref. [15].

In the present work we calculate ground state properties of pure ^4He droplets by means of a variational Monte Carlo (VMC) method. We follow the same procedure as in Ref. [2], but with a different parametrization for the variational wave function. A detailed discussion about the results for the ground state of the droplets is given elsewhere [23]. Here we present

the first microscopic results for the kinetic energy density, which are shown to be in good agreement with the predictions of Ref. [15]. Finally, we use the ground state properties of ^4He droplets as an input in the Feynman-Lekner theory for ^3He impurities. We calculate the binding energy of the ^3He surface states for several ^4He droplets. The results are extrapolated to estimate the binding energy of ^3He atoms on a planar ^4He surface, in good agreement with the experimental value [14].

The work is organized as follows. Section II contains the short description of the VMC method and the results for the kinetic energy density of ^4He droplets. The Feynman-Lekner approach is briefly introduced in Section III, where the results for the Andreev states are also discussed. Section IV is devoted to conclusions.

II. PARTICLE AND KINETIC ENERGY DENSITY OF PURE ^4He DROPLETS

First, we want to calculate the ground state energy, density profile, and kinetic energy density for droplets of given number of particle N . We use the VMC method. In the variational approach a suitable form for the many body wave function $\Psi_v(\mathbf{r}_1, \dots, \mathbf{r}_N)$ is chosen, containing a set of parameters that are varied in order to minimize the energy of the system. The Hamiltonian has the usual form

$$H_0 = \sum_j \frac{-\hbar^2}{2m_4} \nabla_j^2 + \sum_{i < j} v(r_{ij}) \quad , \quad (1)$$

where m_4 is the mass of ^4He atoms. Aziz HFDHE2 potential [24] is used as the pair interatomic potential $v(r_{ij})$.

Several forms of the variational wave function of the pure ^4He liquid have been proposed in the literature [1,2,5,6,23]. They are expected to give very similar results in the context of the present work. We use the formalism of Ref. [23], where the wave function is taken of the same form as in Ref. [2],

$$\Psi_v = \prod_{i \leq N} f_1(r_i) \prod_{i < j \leq N} f_2(r_{ij}) \prod_{i < j < k \leq N} f_3(r_{ij}, r_{jk}, r_{ki}) \quad , \quad (2)$$

but with an improved version of the correlation functions f_1 , f_2 , and f_3 . This choice ensures a correct asymptotic behavior of the wave function far outside the droplet, which has an important role for the ^3He and hydrogen impurity states.

Once the optimal wave function is found, it is used to calculate the one-body density and the kinetic energy density, given by

$$\rho(\mathbf{r}_1) = N \int d\mathbf{r}_2 \dots d\mathbf{r}_N |\Psi_v|^2 \quad , \quad (3)$$

and

$$\tau(\mathbf{r}_1) = \frac{N}{\rho(\mathbf{r}_1)} \int d\mathbf{r}_2 \dots d\mathbf{r}_N \Psi_v^* \left(\frac{-\hbar^2}{2m_4} \nabla_1^2 \right) \Psi_v \quad , \quad (4)$$

respectively.

We consider droplets with 70, 112, 168, 240, 330, and 728 particles. The resulting density profiles and the kinetic energy density are shown in Fig. 1. It is worth noticing that both Monte Carlo [1,2,4–6,23] and density functional [3,11] calculations predict an extrapolated free surface thickness (distance between the points where the density is 90% and 10% of the bulk value) of the order of $6 \div 8 \text{ \AA}$, which is consistent with our results.

The kinetic energy density is shown in the lower part of Fig. 1. It decreases smoothly from the inner value, close to the value of the kinetic energy per particle in bulk liquid, to the asymptotic limit $\mu_4(N)$, i.e., the chemical potential of the drop containing N ^4He atoms. This limit follows from the behavior of the one-body factor which in our parametrization [23] is

$$f_1(r \rightarrow \infty) \propto \sqrt{\rho(r \rightarrow \infty)} \propto \frac{1}{r} \exp \left[- \left(\frac{2m_4 |\mu_4|}{\hbar^2} \right)^{\frac{1}{2}} r \right] \quad , \quad (5)$$

which dominates the $r \rightarrow \infty$ limit of the wave function, yielding

$$\tau(r \rightarrow \infty) = \mu_4 \quad . \quad (6)$$

In Ref. [15] the approximate expression

$$\tau = \tau_0 \left(\frac{\rho}{\rho_0} \right)^n - \frac{\hbar^2}{2m_4} \frac{\nabla^2 \sqrt{\rho}}{\sqrt{\rho}} \quad , \quad (7)$$

was proposed for τ , where τ_0 and ρ_0 are the ground state kinetic energy per particle and the particle density in bulk ^4He at zero pressure, while n is a phenomenological parameter. Equation (7) is an interpolation between the expected behavior of τ in the two opposite limits $\rho \rightarrow \rho_0$ and $\rho \rightarrow 0$. It has never been checked so far with microscopic calculation. In Ref. [15] the values of τ_0 and n were fixed in a phenomenological way to reproduce known properties of ^3He impurities in bulk liquid ^4He , while the saturation density was taken directly from experiments ($\tau_0 = 13.34$ K, $n = 1.76$, $\rho_0 = 0.365\sigma^{-3}$, where $\sigma = 2.556$ Å). The same quantities can be calculated microscopically. For instance, the kinetic energy per particle in bulk liquid at several densities was calculated in Ref. [25] using the variational wave function (2). From those data one extracts $\rho_0 = 0.362\sigma^{-3}$, $\tau_0 = 14.52$ K and $n = 1.77$. In Fig. 2 we compare the VMC results for τ , for droplets with 112 and 728 atoms, with the ones of the approximated formula (7). The solid line corresponds to the variational values of τ_0 , n and ρ_0 , while the dashed line corresponds to the phenomenological parameters of Ref. [15]. In both cases we have used Eq. (7) with the VMC density of the corresponding droplets, and the high frequency statistical fluctuations have been filtered out in the calculation of the second derivative. The figure reveals that the approximate formula (7) works very well, especially for large droplets.

III. LEKNER-FEYNMAN THEORY FOR ^3He IMPURITIES

Once the ground state properties of pure ^4He droplets are obtained, one can calculate the energy and wave function of one ^3He impurity. Consider a droplet consisting of $(N - 1)$ ^4He atoms and one ^3He impurity atom. Since the interatomic potential is the same for ^3He and ^4He , one can write the Hamiltonian in the form

$$H = H_0 + H_I \quad , \quad (8)$$

where H_0 is given in Eq. (1) and

$$H_I = -\frac{\hbar^2}{2m_4} \left(\frac{m_4}{m_3} - 1 \right) \nabla_1^2 \quad . \quad (9)$$

A realistic trial wave function would be

$$\Psi = f(\mathbf{r}_1) \prod_{i=2}^N F(\mathbf{r}_1, \mathbf{r}_i) \Psi_0 \quad , \quad (10)$$

where Ψ_0 is the ground state of the Hamiltonian H_0 , and f and F are variational functions to be determined. In principle, the optimal f and F can be found by minimizing the total energy of the system $\langle \Psi | H | \Psi \rangle$, which can be done either by solving resulting Euler equations or by direct Monte Carlo simulations.

A significant simplification is obtained by taking $F(\mathbf{r}_1, \mathbf{r}_i) \equiv 1$ [26]. This corresponds to the assumption that the correlations between ^3He impurity and ^4He atoms are the same as those among ^4He atoms. The Euler equation for f then becomes

$$-\frac{\hbar^2}{2m_3} \nabla_1 \cdot (\rho(\mathbf{r}_1) \nabla_1 f(\mathbf{r}_1)) + \left(\frac{m_4}{m_3} - 1 \right) f(\mathbf{r}_1) \tau(\mathbf{r}_1) \rho(\mathbf{r}_1) - \lambda f(\mathbf{r}_1) \rho(\mathbf{r}_1) = 0 \quad , \quad (11)$$

where ρ and τ are the particle and kinetic energy densities of the pure ^4He droplet (see Eqs. (3) and (4)). The same equation can be rewritten in the form of a Schrödinger equation

$$-\frac{\hbar^2}{2m_3} \nabla^2 \chi(\mathbf{r}) + V_3(\mathbf{r}) \chi(\mathbf{r}) = \epsilon \chi(\mathbf{r}) \quad , \quad (12)$$

where

$$\chi(\mathbf{r}) = f(\mathbf{r}) \sqrt{\rho(\mathbf{r})} \quad (13)$$

is the impurity wave function, and

$$V_3(\mathbf{r}) = \left(\frac{m_4}{m_3} - 1 \right) \tau(\mathbf{r}) + \frac{\hbar^2}{2m_3} \frac{\nabla^2 \sqrt{\rho(\mathbf{r})}}{\sqrt{\rho(\mathbf{r})}} + \mu_4 \quad (14)$$

is an effective potential seen by the ^3He atom. The chemical potential μ_4 of the droplet of the considered size is added for convenience, so that the eigenvalue ϵ in Eq. (12) is referred to the state in vacuum. In principle, an additional term should appear in the effective potential $V_3(\mathbf{r})$ because the exact ground state of pure ^4He droplet Ψ_0 is approximated by the variational wave function Ψ_v . This term is expected to be small due to the closeness of these two wave functions, and is neglected in the present treatment.

The assumption $F(\mathbf{r}_1, \mathbf{r}_i) \equiv 1$ in Eq. (10) corresponds to the Feynman-Lekner theory [26], already used in the past to predict the properties of the Andreev state on the ^4He free surface [13,15], on films [22], as well as on droplets [10]. Unlike in the previous calculations, where phenomenological ρ and τ were used, here we take them from *ab initio* calculations.

In order to find the energy and the wave function of the impurity on a droplet Eq. (12) is rewritten in spherical coordinates:

$$-\frac{\hbar^2}{2m_3} \frac{d^2}{dr^2} \chi_{nl} + \left(V_3 + \frac{\hbar^2}{2m_3} \frac{l(l+1)}{r^2} \right) \chi_{nl} = \epsilon_{nl} \chi_{nl} \quad . \quad (15)$$

The crucial point is that the effective potential V_3 has a well on the surface of the droplet, so that the lowest eigenstates of the Schrödinger equation are localized on the surface. The potential well originates from a balance between the excess kinetic energy of an ^3He atom in the bulk with respect to the one of ^4He atoms, which tends to push the ^3He atom out, and the He-He interaction, which binds the ^3He atom to the liquid. A typical situation is shown in Fig. 3 for a droplet of 112 atoms. Results for the energy of the lowest ($n = 0$) impurity states on six droplets are given in Fig. 4 as a function of $N^{-1/3}$. We note that a suitable smoothing procedure has been applied to the VMC data in order to avoid spurious effects of statistical fluctuations of ρ in the calculation of the second derivative. This affects mainly the external tail of V_3 where the density is very small. The error bars in Fig. 4 correspond to the consequent inaccuracy in the results, estimated by choosing different smoothing methods. The extrapolation to the case of the planar surface ($N \rightarrow \infty$) can be done by a linear fit, even if the accuracy of the fit is relatively poor. We obtain $\epsilon(\infty) \simeq -4.9$ K, rather close to the experimental estimate $\epsilon(\infty) = (-5.02 \pm 0.03)$ K quoted in Ref. [14].

In Fig. 4 the results of the VMC calculation are compared with predictions of Ref. [10] (full line) obtained using the Feynman-Lekner theory with approximation (7) for τ and density profiles of Ref. [3]. The significant difference between the two predictions is mainly due to the different density profiles. The curvature of the surface profiles in the outer region is underestimated in the density functional calculations of Ref. [3] with respect to the VMC results (see Fig. 5). This makes the potential well for the impurity wider, and the binding

energy lower. The density profiles calculated with a more recent density functional [27,11] are sharper than the ones of Ref. [3] and the corresponding predictions for the ^3He binding energy are closer to the VMC results (dashed lines in Figs. 4 and 5). Part of the remaining discrepancy is due to the use of the approximated expression (7) for τ ; we have checked that the corresponding effect on the ^3He binding energy is small (about 0.1 K).

By solving the Schrödinger equation (12) one finds, above the lowest eigenstate, a spectrum of states with different principal quantum number n and angular momentum l . The general features of the spectrum are the same as in Ref. [10]. In the limit of an infinite droplet the states with different l coincide with those of a 2-dimensional Fermi gas.

To conclude, we stress again the idea of the Feynman-Lekner approach and its limits. The idea is that the two-body correlations between the ^3He impurity and ^4He atoms are taken to be the same as the ones between the ^4He atoms. The ^3He wave function is then expressed by means of the factor $f(\mathbf{r})$ in the many body wave function (10). The form of f is derived variationally, by solving a Schrödinger-like equation. The lowest eigenvalues are localized on the surface of the droplets, as an effect of the different mass of ^3He and ^4He . To improve the Feynman-Lekner theory one should account for the fact that ^3He - ^4He correlations differ from the ^4He - ^4He ones, i.e., one should take $F(\mathbf{r}_1, \mathbf{r}_i) \neq 1$. One possibility is to choose $F = (1 + \phi)$, where ϕ is small. Keeping the terms in the expansion of the total energy up to the second order in ϕ is equivalent to taking into account the energy of an elastic deformation of the residual droplet due to the presence of the impurity. Such corrections have already been studied in Refs. [28] and [20] in the case of liquid ^4He films. The problem of ^3He - ^4He correlations, in the context of variational calculations, has been also discussed in Ref. [29,30] for bulk ^3He - ^4He mixtures. Work in this direction is in progress.

IV. CONCLUSIONS

We present the variational Monte Carlo (VMC) calculation for ^4He droplets with and without ^3He impurities. We have used the variational wave function of Ref. [2] modified as

in Ref. [23]. The results for the density and energy of six droplets, from $N = 70$ to $N = 728$, are close to the predictions of previous calculations. We have calculated the kinetic energy density, for which only approximate estimates have been given so far. Our results show that the analytic formula given in Ref. [15] for the kinetic energy density (Eq. (7)) works very well, and can be safely used in calculations involving the ${}^4\text{He}$ surface.

In the second part of the work we apply the Feynman-Lekner approach to study ${}^3\text{He}$ impurity states on ${}^4\text{He}$ droplets. The impurity wave function turns out to be localized on the surface of the droplets, as expected. The lowest value of the binding energy is almost linearly dependent on $N^{-1/3}$ and the value extrapolated to $N \rightarrow \infty$ is close to the experimental binding energy of the Andreev state on a planar surface. We have also compared our results with the predictions for the ${}^3\text{He}$ binding energies obtained using phenomenological calculations of the density profiles and the kinetic energy density. The general trend of these earlier results is similar to the one of VMC calculations.

REFERENCES

- [1] V.R. Pandharipande, J.G. Zabolitzky, S.C. Pieper, R.B. Wiringa, and U. Helbrecht, Phys. Rev. Lett. **50**, 1676 (1983)
- [2] V.R. Pandharipande, S.C. Pieper, and R.B. Wiringa, Phys. Rev. B **34**, 4571 (1986)
- [3] S. Stringari and J. Treiner, J. Chem. Phys. **87**, 5021 (1987)
- [4] P. Sindzingre, M.L. Klein, and D.M. Ceperley, Phys. Rev. Lett. **63**, 1601 (1989)
- [5] M.V. Rama Krishna and K.B. Whaley, Phys. Rev. Lett. **64**, 1126 (1990); M.V. Rama Krishna and K.B. Whaley, J. Chem. Phys. **93**, 746 (1990); M.V. Rama Krishna and K.B. Whaley, J. Chem. Phys. **93**, 6738 (1990)
- [6] S.A. Chin and E. Krotscheck, Phys. Rev. Lett. **65**, 2658 (1990); S.A. Chin and E. Krotscheck, Phys. Rev. B **45**, 852 (1992)
- [7] A. Scheidemann, J.P. Toennies, and J.A. Northby, Phys. Rev. Lett. **64**, 1899 (1990); A. Scheidemann, B. Schilling and J.P. Toennies, J. Phys. Chem. **97**, 2128 (1993)
- [8] S. Goyal, D.L. Schutt and G. Scoles, Phys. Rev. Lett. **69** 933 (1992); S. Goyal, D.L. Schutt and G. Scoles, Phys. Lett. **196**, 123 (1992); S. Goyal, D.L. Schutt and G. Scoles, J. Phys. Chem. **97**, 2236 (1993)
- [9] R.N. Barnett and K.B. Whaley, J. Chem. Phys. **96**, 2953 (1992); R.N. Barnett and K.B. Whaley (unpublished)
- [10] F. Dalfovo, Z. Phys. D **14**, 263 (1989)
- [11] F. Dalfovo, Z. Phys. D, in press
- [12] A.F. Andreev, Sov. Phys. JETP **23**, 939 (1966)
- [13] W.F. Saam, Phys. Rev. A **4**, 1278 (1971)
- [14] D.O. Edwards and W.F. Saam, in *Progress in Low Temperature Physics*, D.F. Brewer

ed. (North Holland, 1978) Vol. VII A, p.283

- [15] I.B. Mantz and D.O. Edwards, Phys. Rev. B **20**, 4518 (1979)
- [16] F. Dalfovo and S. Stringari, Physica Scripta **38**, 204 (1988)
- [17] N. Pavloff and J. Treiner, J. Low Temp. Phys. **83**, 15 (1991)
- [18] N. Alikacem, R.B. Hallock, R.H. Higley, and D.T. Sprague, J. Low Temp. Phys. **87**, 279 (1992) and references therein
- [19] D.S. Sherrill and D.O. Edwards, Phys. Rev. B **31**, 1338 (1985)
- [20] E. Krotscheck, M. Saarela, and J.L. Epstein, Phys. Rev. B **38**, 111 (1988)
- [21] N. Pavloff and J. Treiner, J. Low Temp. Phys. **83**, 331 (1991)
- [22] J. Treiner, J. Low Temp. Phys **92**, 1 (1993)
- [23] A. Belić and S.C. Pieper, ICTP preprint IC/94/2, submitted to Phys. Rev. B.
- [24] R.A. Aziz, V.P.S. Nain, J.S. Carley, W.L. Taylor, and G.T. McConville, J. Chem. Phys. **70**, 4330 (1979).
- [25] Q.N. Usmani, S. Fantoni, and V.R. Pandharipande, Phys. Rev. B **26**, 6123 (1982)
- [26] J. Lekner, Phil. Mag. **22**, 669 (1970)
- [27] J. Dupont-Roc, M. Himbert, N. Pavloff, and J. Treiner, J. Low Temp. Phys **81**, 31 (1990)
- [28] R.A. Guyer, M.D. Miller, and J. Yaple, Phys. Rev. B **25**, 4570 (1982)
- [29] J. Boronat, A. Fabrocini, and A. Polls, J. Low Temp. Phys. **91**, 275 (1993)
- [30] E. Krotscheck and M. Saarela, Phys. Rep. **232**, 1 (1993)

FIGURES

FIG. 1. Particle density and kinetic energy density for droplets of 70 (squares), 112 (plus signs), 168 (circles), 240 (crosses), 330 (triangles), and 728 (diamonds) atoms.

FIG. 2. Kinetic energy density for droplets of 728 and 112 atoms. Circles: VMC; solid lines: Eq.(7) with $\rho_0 = 0.362\sigma^{-3}$, $\tau_0 = 14.52$ K and $n = 1.77$; dashed lines: the same equation with $\rho_0 = 0.365\sigma^{-3}$, $\tau_0 = 13.34$ K and $n = 1.76$.

FIG. 3. Lowest eigenstate of one ^3He impurity on a droplet of 112 atoms. Solid line: ^3He wave function; squares: ^4He density; circles: effective potential V_3 .

FIG. 4. Binding energy of the lowest impurity state as a function of $N^{-1/3}$. Points with error bars: Feynman-Lekner theory with τ and ρ from VMC calculations; solid line: Feynman-Lekner results of Ref. [10]; dashed line: same theory but with more recent density functional calculations for the density profiles, as in Ref.[11].

FIG. 5. Surface profile for a droplet of 112 atoms. Points: VMC results; solid line: density functional results of Ref. [3]; dashed line: density functional results of Ref.[11].